


Iridodialysis

Gilberto Krasilchik¹ , Nicole Bulgarão Maricondi Almeida² , Newton Kara-Junior² 

1. Ophthalmology Department, Faculdade de Medicina, Santa Casa de Misericórdia de São Paulo, São Paulo, SP, Brazil.

2. Ophthalmology Department, Hospital das Clínicas, Universidade de São Paulo, São Paulo, SP, Brazil.

Iridodialysis refers to the separation of the iris from its attachment to the ciliary body. This condition commonly arises due to trauma or complications during intraocular surgery, given the inherently weak structure of the iris root⁽¹⁾. The iris plays a crucial role in controlling pupil diameter⁽¹⁾. Consequently, iris defects can lead not only to cosmetic deformations, such as polycoria and ectopic pupil, but also to significant functional impairments. These include monocular diplopia, photophobia⁽²⁾, reduced visual quality, severe glare, and decreased contrast sensitivity⁽¹⁾.



REFERENCES

1. Ferro Desideri L, Arun K, Doherty G, Bernardi E, Anguita R. Iris reconstruction: A surgeon's guide. *J Clin Med.* 2024;13(9):2706.
2. Wan W, Shi L, Li C. Comparing safety and efficiency of two closed-chamber techniques for iridodialysis repair - a retrospective clinical study. *BMC Ophthalmol.* 2018;18(1):311.

Submitted for publication: March 10, 2025

Accepted for publication: March 14, 2025

Funding: This study received no specific financial support.

Disclosure of potential conflicts of interest: The authors declare no potential conflicts of interest.

Corresponding author: Nicole B.M. Almeida.

E-mail: nickbma204@gmail.com

Informed consent was obtained from all patients included in this study.

Edited by

Editor-in-Chief: Newton Kara-Júnior

Authors' information

NKJr is Editor-in-Chief of ABO. This article was independently handled by a member of the Editorial Board.

 This content is licensed under a Creative Commons Attributions 4.0 International License.